

Euler-Mahonian polynomials for $C_a \wr S_n$

Dan Bernstein

Department of Mathematics
The Weizmann Institute of Science
Rehovot 76100, Israel
`dan.bernstein@weizmann.ac.il`

February 1, 2008

1 Introduction

Let S_n denote the symmetric group on $\{1, \dots, n\}$. The classical *descent number* and *major index* statistics on S_n are defined by

$$\text{des}(\sigma) := \sum_{i=1}^{n-1} \chi(\sigma(i) > \sigma(i+1))$$

and

$$\text{maj}(\sigma) := \sum_{i=1}^{n-1} i \chi(\sigma(i) > \sigma(i+1))$$

respectively, where $\sigma \in S_n$ and $\chi(p)$ equals 1 if the statement p is true and 0 if p is false.

The classical Eulerian polynomials $A_n(t)$ may be defined by

$$A_n(t) := \sum_{\sigma \in S_n} t^{\text{des}(\sigma)}$$

and their “maj” q -analogues, also known as the q -maj *Euler-Mahonian polynomials*, may be defined by

$$\text{maj} A_n(t, q) := \sum_{\sigma \in S_n} t^{\text{des}(\sigma)} q^{\text{maj}(\sigma)}. \quad (1)$$

Carlitz [Car75] has given a recurrence formula for the coefficients of ${}^{\text{maj}}A_n(t, q)$: with ${}^{\text{maj}}A_n(t, q) = \sum_{s \geq 0} t^s {}^{\text{maj}}A_{n,s}(q)$,

$${}^{\text{maj}}A_{n,s}(q) = (1 + q + \cdots + q^s) {}^{\text{maj}}A_{n-1,s}(q) + (q^s + q^{s+1} + \cdots + q^{n-1}) {}^{\text{maj}}A_{n-1,s-1}(q). \quad (2)$$

Gessel [Ges77] has obtained the exponential generating function for the quotients $\frac{{}^{\text{maj}}A_n(t, q)}{(1-t)(1-qt)\cdots(1-q^n t)}$,

$$\sum_{n \geq 0} \frac{u^n}{n!} \frac{{}^{\text{maj}}A_n(t, q)}{(1-t)(1-qt)\cdots(1-q^n t)} = \sum_{s \geq 0} t^s e^{u(1+q+\cdots+q^s)}. \quad (3)$$

Identity (3) is known as Carlitz's identity.

Let C_a be the cyclic group of order a , and let $C_a \wr S_n$ be its wreath product with the symmetric group S_n , which comprises *colored permutations*.

The problem of extending the distribution of (des, maj) to the hyperoctahedral group $B_n = C_2 \wr S_n$ was first suggested by Foata. Adin, Brenti and Roichman [ABR01] have given a solution to Foata's problem in the form of two pairs of statistics, $(\text{ndes}, \text{nmaj})$ and $(\text{fdes}, \text{fmaj})$. Later, Biagioli [Bia03] has given a generalization of Carlitz's identity to the even-signed permutation group D_n . More recently, Bagno [Bag04] has extended the des and maj statistics to the wreath products $C_a \wr S_n$ in two different ways, $(\text{ndes}, \text{nmaj})$ and $(\text{lides}, \text{lmaj})$, giving two further generalizations of Carlitz's formula.

In a recent paper [RR04], Regev and Roichman introduced the order $<_L$ and the $C_a \wr S_n$ statistics des_L (the L -descent number) and $\overleftarrow{\min}_L$ (number of L -colored right-to-left minima), and studied the distribution of $\overleftarrow{\min}_L$ on $C_a \wr S_n$ and on the subset $\{\sigma \in C_a \wr S_n : \overleftarrow{\min}_L(\sigma) = \text{des}_L(\sigma)\}$.

Here we define the $\text{rmaj}_{L,n}$ (L -reverse major index) statistic on $C_a \wr S_n$ and study the distribution of des_L and the bi-statistic $(\text{des}_L, \text{rmaj}_{L,n})$. We obtain new wreath-product analogues of the Eulerian and q -Euler-Mahonian polynomials, and a generalization of Carlitz's identity (see Corollary 4.5).

2 Preliminaries

2.1 The Group $C_a \wr S_n$

Let C_a be the (multiplicative) cyclic group of order a : $\alpha := e^{\frac{2\pi i}{a}}$ and $C_a := \{\alpha^t : 0 \leq t \leq a-1\}$, and let $C_a \wr S_n$ be its wreath product with S_n .

Elements of $C_a \wr S_n$ can be regarded as *indexed permutations* or *colored permutations*—those permutations σ of the set $\{\alpha^t i : 0 \leq t \leq a-1, 1 \leq i \leq n\}$ satisfying

$$\sigma(\beta j) = \beta \sigma(j) \quad \forall \beta \in C_a, 1 \leq j \leq n.$$

We shall write colored permutations using the *window notation*

$$\sigma = [\sigma(1), \dots, \sigma(n)]$$

and denote

$$|\sigma| := [|\sigma(1)|, \dots, |\sigma(n)|] \in S_n.$$

Note that S_n is a subgroup of $C_a \wr S_n$. $C_a \wr S_n$ is a Coxeter group if and only if $a = 1$ or 2 . In particular, for $a = 2$, $C_2 \wr S_n = B_n$ is the hyperoctahedral group, whose elements are the *signed permutations*.

2.2 q -analogues

Definition 2.1. For an integer $n \geq 1$, the q -analogue of n is

$$[n]_q := 1 + q + \dots + q^{n-1}.$$

Definition 2.2. For an integer $n \geq 0$, define

$$(\alpha; q)_n := \begin{cases} 1, & \text{if } n = 0; \\ (1 - \alpha)(1 - \alpha q) \cdots (1 - \alpha q^{n-1}), & \text{if } n > 0. \end{cases}$$

3 Statistics on $C_a \wr S_n$

In this section we present Regev and Roichman's $<_L$ order and various statistics based on it.

Definition 3.1 ([RR04, Definition 4.4]). A subset $L \subseteq \{0, 1, \dots, a-1\}$ determines a linear order $<_L$ on $\{\alpha^v j : 0 \leq v \leq a-1, 0 \leq j \leq n\} \cup \{0\}$ as follows:

Let $U = \{0, \dots, a-1\} \setminus L$ be the complement of L in $\{0, \dots, a-1\}$.

If $v \in L$ then $\alpha^v j <_L 0$ for every $1 \leq j \leq n$. If $v \in U$ then $\alpha^v j >_L 0$ for every $1 \leq j \leq n$.

For $u, v \in L$ (resp. $\in U$) (not necessarily distinct) and $i \neq j \in [n]$, $\alpha^v i <_L \alpha^u j$ if and only if $i > j$ (resp. $i < j$).

Then, for each $1 \leq j \leq n$, order each subset $\{\alpha^v j : v \in L\}$ (and each subset $\{\alpha^v j : v \in U\}$) in an arbitrary linear order.

Example 3.2. Let $a = 4$ and $L = \{2, 3\}$, then $U = \{0, 1\}$. We can choose the following order

$$\begin{aligned} \alpha^2 n <_L \alpha^3 n <_L \alpha^2(n-1) <_L \alpha^3(n-1) <_L \cdots <_L \alpha^2 <_L \alpha^3 <_L 0 \\ 0 <_L \alpha <_L 1 <_L \alpha 2 <_L \cdots <_L \alpha(n-1) <_L (n-1) <_L \alpha n <_L n. \end{aligned}$$

Definition 3.3 ([RR04, Definition 4.6]). Let $L \subseteq \{0, 1, \dots, a-1\}$.

1. The L -descent set of $\sigma \in C_a \wr S_n$ is

$$\text{Des}_L(\sigma) := \{0 \leq i \leq n-1 : \sigma(i) >_L \sigma(i+1)\}$$

where $\sigma(0) := 0$.

2. The L -descent number is

$$\text{des}_L(\sigma) := |\text{Des}_L(\sigma)|.$$

Definition 3.4. The L -reverse major index of $\sigma \in C_a \wr S_n$ is

$$\text{rmaj}_{L,n}(\sigma) := \sum_{i \in \text{des}_L(\sigma)} n - i.$$

Remark 3.5. In the $a = 2$ case, the descent set is often defined as

$$\widetilde{\text{Des}}_L(\sigma) := \{1 \leq i \leq n : \sigma(i) >_L \sigma(i+1)\}$$

where $\sigma(n+1) := 0$ (see for example [FH97]). It is easy to see that $\widetilde{\text{Des}}_L(\sigma) = \{n-i : i \in \text{Des}_{\{1, \dots, n\} \setminus L}(\sigma \cdot [n, n-1, \dots, 1])\}$, so $\widetilde{\text{des}}_L(\sigma) := |\{\widetilde{\text{Des}}_L(\sigma)\}| = \text{des}_{\{1, \dots, n\} \setminus L}(\sigma \cdot [n, \dots, 1])$ and $\widetilde{\text{maj}}_L(\sigma) := \sum_{i \in \widetilde{\text{des}}_L(\sigma)} i = \text{rmaj}_{L,n}(\sigma \cdot [n, \dots, 1])$. Since multiplication by $[n, \dots, 1]$ is an involution of $C_a \wr S_n$, we get that the bi-statistics $(\widetilde{\text{des}}_L, \widetilde{\text{maj}}_L)$ and $(\text{des}_{\{1, \dots, n\} \setminus L}, \text{rmaj}_{\{1, \dots, n\} \setminus L, n})$ have the same distribution on $C_a \wr S_n$. Thus the results in the following sections can be easily adapted to the “tilde” statistics.

Remark 3.6. For $\sigma \in S_n$, let $\text{rmaj}_n(\sigma) := \text{rmaj}_{\emptyset, n}(\sigma)$. A bijective argument shows that maj and rmaj_n are equidistributed on $\{\sigma \in S_n : \text{des}(\sigma) = s\}$ for every s . Thus in (1), maj can be replaced by rmaj_n .

Definition 3.7. Define $\phi_n : C_a \wr S_{n-1} \times \{0, \dots, n-1\} \times \{0, \dots, a-1\} \rightarrow C_a \wr S_n$ by

$$\phi_n(\sigma, r, t) := [\sigma_1, \dots, \sigma_r, \alpha^t n, \sigma_{r+1}, \dots, \sigma_{n-1}]$$

where $\sigma = [\sigma_1, \dots, \sigma_{n-1}]$.

It is easy to see that ϕ_n is a bijection.

Lemma 3.8. Let $\sigma = [\sigma_1, \dots, \sigma_{n-1}] \in C_a \wr S_{n-1}$, $L \subseteq \{0, \dots, a-1\}$ and $\text{Des}_L(\sigma) = \{i_1, \dots, i_s\}$, $i_1 < \dots < i_s$ ($s = \text{des}_L(\sigma)$). Let $\{i_{s+1}, \dots, i_n\} = \{0, \dots, n-1\} \setminus \text{Des}_L(\sigma)$, $n-1 = i_{s+1} > \dots > i_n$ (i.e. the non-descents of σ , from right to left). Then for $1 \leq k \leq n$ and $t \in \{0, \dots, a-1\}$,

$$\text{des}_L(\phi_n(\sigma, i_k, t)) = \begin{cases} s, & \text{if } k < s+1 \text{ or } k = s+1, t \notin L; \\ s+1, & \text{if } k > s+1 \text{ or } k = s+1, t \in L. \end{cases}$$

and

$$\text{rmaj}_{L,n}(\phi_n(\sigma, i_k, t)) = \begin{cases} \text{rmaj}_{L,n-1}(\sigma) + k, & \text{if } t \in L; \\ \text{rmaj}_{L,n-1}(\sigma) + k - 1, & \text{if } t \notin L, \end{cases}$$

Proof. We consider the three possible cases:

Case 1. $1 \leq k \leq s$. In this case,

$$\tilde{\sigma} := \phi_n(\sigma, i_k, t) = [\sigma_1, \dots, \sigma_{i_1}, \dots, \sigma_{i_k}, \alpha^t n, \sigma_{i_k+1}, \dots, \sigma_{i_{k+1}}, \dots, \sigma_{i_s}, \dots, \sigma_{n-1}],$$

thus the descents to the right of σ_{i_k+1} are shifted one position to the right, and the $k-1$ descents to the left of σ_{i_k} remain in place. If $t \in L$, then $\sigma_{i_k} >_L \alpha^t n <_L \sigma_{i_k+1}$, so the descent at i_k is also shifted one position to the right. If $t \notin L$, then $\sigma_{i_k} <_L \alpha^t n >_L \sigma_{i_k+1}$, leaving the descent at i_k . The contribution to $\text{rmaj}_{L,n}(\tilde{\sigma})$ of each descent shifted one position to the right is the same as its contribution to $\text{rmaj}_{L,n-1}(\sigma)$, whereas the contribution to $\text{rmaj}_{L,n}(\tilde{\sigma})$ of each descent left in place is 1 more than its contribution to $\text{rmaj}_{L,n-1}(\sigma)$.

Case 2. $k = s+1$. In this case,

$$\tilde{\sigma} := \phi_n(\sigma, n-1, t) = [\sigma_1, \dots, \sigma_{n-1}, \alpha^t n],$$

thus all $s = k-1$ descents remain in place, each contributing 1 more to $\text{rmaj}_{L,n}(\tilde{\sigma})$ than to $\text{rmaj}_{L,n-1}(\sigma)$. If and only if $t \in L$, $\sigma_{n-1} >_L \alpha^t n$ so there is an additional descent at $n-1$, which contributes 1 to $\text{rmaj}_{L,n}(\tilde{\sigma})$.

Case 3. $s + 1 < k \leq n$. In this case,

$$\tilde{\sigma} := \phi_n(\sigma, i_k, t) = [\sigma_1, \dots, \sigma_{i_r}, \dots, \sigma_{i_k}, \alpha^t n, \sigma_{i_k+1}, \dots, \sigma_{i_{r+1}}, \dots, \sigma_{n-1}]$$

where r is the number of descents to the left of σ_{i_k} , whence

$$i_k = (n - 1) - (k - (s + 1)) - (s - r) = n - k + r.$$

In this case the descents to the right of σ_{i_k} are shifted one position to the right, and the r descents to the left of σ_{i_k} remain in place. The contribution to $\text{rmaj}_{L,n}(\tilde{\sigma})$ of each descent shifted one position to the right is the same as its contribution to $\text{rmaj}_{L,n-1}(\sigma)$, whereas the contribution to $\text{rmaj}_{L,n}(\tilde{\sigma})$ of each descent left in place is 1 more than its contribution to $\text{rmaj}_{L,n-1}(\sigma)$. If $t \in L$, then there is an additional descent at $i_k = n - k + r$, whose contribution to $\text{rmaj}_{L,n}(\tilde{\sigma})$ is $k - r$. If $t \notin L$, then the additional descent is at $i_k + 1 = n - k + r + 1$, contributing $k - r - 1$ to $\text{rmaj}_{L,n}(\tilde{\sigma})$. $\square$

4 $C_a \wr S_n$ q -maj Euler-Mahonian Polynomials

In this section we define q -maj Euler-Mahonian polynomials for $C_a \wr S_n$ and give generalizations of the results by Carlitz and Gessel.

For $L \subseteq \{0, \dots, a - 1\}$, let ${}^{\text{maj}}A_{a,L,n}(t, q)$ be the generating polynomial for $C_a \wr S_n$ by the bi-statistic $(\text{des}_L, \text{rmaj}_{L,n})$, i.e.

$${}^{\text{maj}}A_{a,L,n}(t, q) := \sum_{\sigma \in C_a \wr S_n} t^{\text{des}_L(\sigma)} q^{\text{rmaj}_{L,n}(\sigma)}.$$

Remark 4.1. For $n = 1$,

$${}^{\text{maj}}A_{a,L,1}(t, q) = \sum_{t=0}^{a-1} t^{\text{des}_L([\alpha^t])} q^{\text{rmaj}_{L,1}([\alpha^t])} = \ell t q + (a - \ell)$$

where $\ell = |L|$ depends only on the number of elements in L and not on the choice of elements.

The following is a generalization of (2).

Proposition 4.2. *With ${}^{\text{maj}}A_{a,L,n}(t, q) = \sum_{s \geq 0} t^s {}^{\text{maj}}A_{a,L,n,s}(q)$, the coefficients ${}^{\text{maj}}A_{a,L,n,s}(q)$ satisfy the recurrence*

$${}^{\text{maj}}A_{a,L,n,s}(q) = (a[s+1]_q - \ell) {}^{\text{maj}}A_{a,L,n-1,s}(q) + (aq^s[n-s]_q + \ell q^n) {}^{\text{maj}}A_{a,L,n-1,s-1}(q) \quad (4)$$

where $\ell = |L|$.

Proof. By definition,

$${}^{\text{maj}}A_{a,L,n,s}(q) = \sum_{\substack{\tilde{\sigma} \in C_a \wr S_n \\ \text{des}_L(\tilde{\sigma}) = s}} q^{\text{rmaj}_{L,n}(\tilde{\sigma})}.$$

By the bijectivity of ϕ_n and Lemma 3.8,

$$\{\tilde{\sigma} \in C_a \wr S_n : \text{des}_L(\tilde{\sigma}) = s\} = \phi_n(A \uplus B \uplus C \uplus D)$$

where $\uplus$ denotes disjoint union and

$$A := \{(\sigma, i_k, t) : \sigma \in C_a \wr S_{n-1}, \text{Des}_L(\sigma) = \{i_1, \dots, i_s\}, \ 1 \leq k \leq s, \ t \in L\}$$

$$B := \{(\sigma, i_k, t) : \sigma \in C_a \wr S_{n-1}, \{0, \dots, n-1\} \setminus \text{Des}_L(\sigma) = \{i_s, \dots, i_n\}, \ s \leq k \leq n, \ t \in L\}$$

$$C := \{(\sigma, i_k, t) : \sigma \in C_a \wr S_{n-1}, \text{Des}_L(\sigma) \uplus \{n-1\} = \{i_1, \dots, i_{s+1}\}, \ 1 \leq k \leq s+1, \ t \notin L\}$$

$$D := \{(\sigma, i_k, t) : \sigma \in C_a \wr S_{n-1}, \{0, \dots, n-2\} \setminus \text{Des}_L(\sigma) = \{i_{s+1}, \dots, i_n\}, \ s+1 \leq k \leq n, \ t \notin L\}.$$

Note that in the definition of A and C , $\text{des}_L(\sigma) = s$, whereas in the definition of B and D , $\text{des}_L(\sigma) = s-1$.

By the second part of Lemma 3.8,

$$\begin{aligned}
\sum_{\substack{\sigma \in C_a \wr S_n \\ \text{des}_L(\sigma) = s}} q^{\text{rmaj}_{L,n}(\sigma)} &= \sum_{(\sigma, i_k, t) \in A} q^{\text{rmaj}_{L,n}(\phi_n(\sigma, i_k, t))} + \sum_{(\sigma, i_k, t) \in B} q^{\text{rmaj}_{L,n}(\phi_n(\sigma, i_k, t))} \\
&+ \sum_{(\sigma, i_k, t) \in C} q^{\text{rmaj}_{L,n}(\phi_n(\sigma, i_k, t))} + \sum_{(\sigma, i_k, t) \in D} q^{\text{rmaj}_{L,n}(\phi_n(\sigma, i_k, t))} \\
&= \sum_{(\sigma, i_k, t) \in A} q^{\text{rmaj}_{L,n-1}(\sigma) + k} + \sum_{(\sigma, i_k, t) \in B} q^{\text{rmaj}_{L,n-1}(\sigma) + k} \\
&+ \sum_{(\sigma, i_k, t) \in C} q^{\text{rmaj}_{L,n-1}(\sigma) + k - 1} + \sum_{(\sigma, i_k, t) \in D} q^{\text{rmaj}_{L,n-1}(\sigma) + k - 1} \\
&= \ell q[s]_q \sum_{\substack{\sigma \in C_a \wr S_{n-1} \\ \text{des}_L(\sigma) = s}} q^{\text{rmaj}_{L,n-1}(\sigma)} \\
&+ \ell q^s[n - s + 1]_q \sum_{\substack{\sigma \in C_a \wr S_{n-1} \\ \text{des}_L(\sigma) = s-1}} q^{\text{rmaj}_{L,n-1}(\sigma)} \\
&+ (a - \ell)[s + 1]_q \sum_{\substack{\sigma \in C_a \wr S_{n-1} \\ \text{des}_L(\sigma) = s}} q^{\text{rmaj}_{L,n-1}(\sigma)} \\
&+ (a - \ell)q^s[n - s]_q \sum_{\substack{\sigma \in C_a \wr S_{n-1} \\ \text{des}_L(\sigma) = s-1}} q^{\text{rmaj}_{L,n-1}(\sigma)} \\
&= (a[s + 1]_q - \ell)^{\text{maj}} A_{a,L,n-1,s} \\
&+ (aq^s[n - s]_q + \ell q^n)^{\text{maj}} A_{a,L,n-1,s-1}.
\end{aligned}$$

□

By Proposition 4.2 and Remark 4.1, $^{\text{maj}}A_{a,L,n}(t, q)$ does not depend on the choice of elements in L but only on their number $\ell = |L|$. Therefore the polynomials

$$^{\text{maj}}A_{a,\ell,n}(t, q) := ^{\text{maj}}A_{a,L,n}(t, q) \quad |L| = \ell$$

and

$$^{\text{maj}}A_{a,\ell,n,s}(q) := ^{\text{maj}}A_{a,L,n,s}(q) \quad |L| = \ell$$

are well-defined.

The following lemma is a generalization of [FH04, equation (10.3)].

Lemma 4.3. *For every three integers $n > 0$, $a > 0$, $0 \leq \ell \leq a$,*

$$(1 - q)^{\text{maj}} A_{a,\ell,n}(t, q) = (a - (1 - q)\ell)(1 - tq^n)^{\text{maj}} A_{a,\ell,n-1}(t, q) - aq(1 - t)^{\text{maj}} A_{a,\ell,n-1}(tq, q). \quad (5)$$

Proof. Multiply both sides of (4) by $(1 - q)t^s$ and sum over all $-\infty < s < \infty$ to get

$$\begin{aligned} & (1 - q)^{\text{maj}} A_{a,\ell,n}(t, q) \\ &= \sum_{s=-\infty}^{\infty} (a(1 - q^{s+1}) - (1 - q)\ell)t^s \text{maj} A_{a,\ell,n-1,s}(q) \\ & \quad + \sum_{s=-\infty}^{\infty} (aq^s(1 - q^{n-s}) + (1 - q)\ell q^n)t^s \text{maj} A_{a,\ell,n-1,s-1}(q) \\ &= \sum_{s=-\infty}^{\infty} ((a - (1 - q)\ell)t^s - aq(qt)^s) \text{maj} A_{a,\ell,n-1,s}(q) \\ & \quad + \sum_{s=-\infty}^{\infty} t((-aq^n + (1 - q)\ell q^n)t^s + a(qt)^s) \text{maj} A_{a,\ell,n-1,s}(q) \\ &= \sum_{s=-\infty}^{\infty} ((a - (1 - q)\ell)(1 - tq^n)t^s - aq(1 - t)(qt)^s) \text{maj} A_{a,\ell,n-1,s}(q) \\ &= (a - (1 - q)\ell)(1 - tq^n)^{\text{maj}} A_{a,\ell,n-1}(t, q) - aq(1 - t)^{\text{maj}} A_{a,\ell,n-1}(tq, q). \quad \square \end{aligned}$$

Proposition 4.4. *For every three integers $n > 0$, $a > 0$, $0 \leq \ell \leq a$,*

$$\frac{\text{maj} A_{a,\ell,n}(t, q)}{(t; q)_{n+1}} = \sum_{s \geq 0} t^s (a[s + 1]_q - \ell)^n. \quad (6)$$

Proof. By induction on n . For $n = 1$,

$$\frac{\text{maj} A_{a,\ell,1}(t, q)}{(t; q)_2} = \frac{a - \ell(1 - tq)}{(1 - t)(1 - tq)} = \sum_{s \geq 0} t^s (a[s + 1]_q - \ell).$$

For $n > 1$, divide both sides of (5) by $(1 - q)(t; q)_{n+1}$ and use the induction

hypothesis to get

$$\begin{aligned}
& \frac{{}^{\text{maj}}A_{a,\ell,n}(t, q)}{(t; q)_{n+1}} \\
&= \frac{a - (1 - q)\ell}{(1 - q)(t; q)_n} {}^{\text{maj}}A_{a,\ell,n-1}(t, q) - \frac{aq(1 - t)}{(1 - q)(t; q)_{n+1}} {}^{\text{maj}}A_{a,\ell,n-1}(tq, q) \\
&= \left(\frac{a}{1 - q} - \ell \right) \sum_{s \geq 0} t^s (a[s + 1]_q - \ell)^{n-1} - \frac{aq}{1 - q} \sum_{s \geq 0} t^s q^s (a[s + 1]_q - \ell)^{n-1} \\
&= \sum_{s \geq 0} t^s \left(\frac{a}{1 - q} - \ell - \frac{aq^{s+1}}{1 - q} \right) (a[s + 1]_q - \ell)^{n-1} \\
&= \sum_{s \geq 0} t^s (a[s + 1]_q - \ell)^n.
\end{aligned}$$

□

As a corollary, we get a generalization of (3).

Corollary 4.5. *For every $a > 0$, $0 \leq \ell \leq a$,*

$$\sum_{n \geq 0} \frac{u^n}{n!} \frac{{}^{\text{maj}}A_{a,\ell,n}(t, q)}{(t; q)_{n+1}} = \sum_{s \geq 0} t^s e^{u(a[s+1]_q - \ell)}. \quad (7)$$

5 $C_a \wr S_n$ Eulerian Polynomials

For $L \subseteq \{0, \dots, a - 1\}$, let $A_{a,L,n}(t)$ be the generating polynomial for $C_a \wr S_n$ by the statistic des_L , i.e.

$$A_{a,L,n}(t) := \sum_{\sigma \in C_a \wr S_n} t^{\text{des}_L(\sigma)}.$$

Clearly $A_{a,L,n}(t) = {}^{\text{maj}}A_{a,L,n}(t, 1)$ and the polynomials

$$A_{a,\ell,n}(t) := A_{a,L,n}(t) \quad |L| = \ell$$

are well-defined.

Proposition 5.1. *1. With $A_{a,\ell,n}(t) = \sum_{s \geq 0} t^s A_{a,\ell,n,s}$, the coefficients $A_{a,\ell,n,s}$ satisfy the recurrence*

$$A_{a,\ell,n,s} = (a(s + 1) - \ell) A_{a,\ell,n-1,s} + (a(n - s) + \ell) A_{a,\ell,n-1,s-1}. \quad (8)$$

2.

$$A_{n,a,\ell}(t) = (a - \ell + (a(n-1) + \ell)t)A_{a,\ell,n-1}(t) + at(1-t)A'_{a,\ell,n-1}(t). \quad (9)$$

3.

$$\frac{A_{a,\ell,n}(t)}{(1-t)^{n+1}} = \sum_{s \geq 0} t^s (a(s+1) - \ell)^n. \quad (10)$$

4.

$$\sum_{n \geq 0} \frac{u^n}{n!} \frac{A_{a,\ell,n}(t)}{(1-t)^{n+1}} = \sum_{s \geq 0} t^s e^{u(a(s+1) - \ell)} = \frac{e^{(a-\ell)u}}{1 - te^{au}}. \quad (11)$$

5.

$$\sum_{n \geq 0} \frac{u^n}{n!} A_{a,\ell,n}(t) = \frac{1-t}{-te^{\ell(1-t)u} + e^{(a-\ell)u(t-1)}}. \quad (12)$$

Proof. (8), (10) and (11) follow from (4), (6) and (7) respectively by setting $q = 1$. (12) follows from (11) by substituting $(1-t)u$ for u and multiplying both sides by $1 = t$. Finally, to see (9), multiply both sides of (8) by t^s and take the sum over all $s \geq 0$ to get

$$\begin{aligned} A_{a,\ell,n}(t) &= \sum_{s \geq 0} (a(s+1) - \ell)t^s A_{a,\ell,n-1,s} + \sum_{s \geq 0} (a(n-s) + \ell)t^s A_{a,\ell,n-1,s-1} \\ &= \sum_{s \geq 0} (a(s+1) - \ell)t^s A_{a,\ell,n-1,s} + t \sum_{s \geq 0} (a(n-s-1) + \ell)t^s A_{a,\ell,n-1,s} \\ &= \sum_{s \geq 0} (a - \ell + (a(n-1) + \ell)t)t^s A_{a,\ell,n-1,s} \\ &\quad + t \sum_{s \geq 0} a(1-t)st^{s-1} A_{a,\ell,n-1,s} \\ &= (a - \ell + (a(n-1) + \ell)t)A_{a,\ell,n-1}(t) + at(1-t)A'_{a,\ell,n-1}(t). \end{aligned}$$

□

References

- [ABR01] Ron M. Adin, Francesco Brenti, and Yuval Roichman. Descent numbers and major indices for the hyperoctahedral group. *Adv. in Appl. Math.*, 27(2-3):210–224, 2001.

- [Bag04] Eli Bagno. Euler-Mahonian parameters on colored permutation groups. <http://www.math.biu.ac.il/~bagnoe/majorver2.pdf>, 2004.
- [Bia03] Riccardo Biagioli. Major and descent statistics for the even-signed permutation group. *Adv. in Appl. Math.*, 31:163–179, 2003.
- [Car75] Leonard Carlitz. A combinatorial property of q -Eulerian numbers. *Amer. Math. Monthly*, 82:51–54, January 1975.
- [FH97] Dominique Foata and Guo-Niu Han. Calcul basique des permutations signées. I. Longueur et nombre d’inversions. *Adv. in Appl. Math.*, 18(4):489–509, 1997.
- [FH04] Dominique Foata and Guo-Niu Han. The q -series in combinatorics; permutation statistics. Preliminary version, 2004.
- [Ges77] Ira M. Gessel. *Generating functions and Enumeration of Sequences*. PhD thesis, M.I.T., 1977.
- [RR04] Amitai Regev and Yuval Roichman. Statistics on wreath products and generalized binomial-Stirling numbers. [arXiv:math.CO/0404354](https://arxiv.org/abs/math.CO/0404354), 2004.